

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

SCHEDULING STATUS:

S2

PROPRIETARY NAME AND DOSAGE FORM:

TEXA ALLERGY SYRUP syrup

Read all of this leaflet carefully because it contains important information for you.

TEXA ALLERGY SYRUP is available without a doctor's prescription, for you to treat a mild illness. Nevertheless, you still need to use **TEXA ALLERGY SYRUP** carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Do not share **TEXA ALLERGY SYRUP** with any other person.
- Ask your pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve after 7 days.

1. WHAT TEXA ALLERGY SYRUP CONTAINS:

The active substance is cetirizine dihydrochloride.

One ml of **TEXA ALLERGY SYRUP** contains 1 mg cetirizine dihydrochloride.

The other ingredients are acetic acid, banana flavour, glycerol, methyl parahydroxybenzoate, propylene glycol, propyl hydroxybenzoate, purified water, saccharin sodium, sodium acetate and sorbitol 70 %.

TEXA ALLERGY SYRUP contains the preservatives methyl parahydroxybenzoate and propyl hydroxybenzoate. **TEXA ALLERGY SYRUP** also contains sweeteners (glycerol 200 mg/ml, saccharin sodium 1 mg/ml and sorbitol 70 % solution 450 mg/ml) (see **Important information about some of the ingredients of TEXA ALLERGY SYRUP**).

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

2. WHAT TEXA ALLERGY SYRUP IS USED FOR:

TEXA ALLERGY SYRUP belongs to a group of medicines called long-acting non-sedating antihistamines.

TEXA ALLERGY SYRUP is used for the relief of allergic conditions such as:

- allergic rhinitis (allergy related stuffy or runny nose, sneezing) and hay fever
- allergic skin conditions such as urticaria (rash of round, red weals on the skin which itch intensely, sometimes with dangerous swelling, caused by an allergic reaction).

3. BEFORE YOU TAKE TEXA ALLERGY SYRUP:

Do not take TEXA ALLERGY SYRUP:

- if you are hypersensitive (allergic) to cetirizine, hydroxyzine, piperazine or any other antihistamines, or to any of the ingredients of **TEXA ALLERGY SYRUP** (see **WHAT TEXA ALLERGY SYRUP CONTAINS**)
- if you have severe kidney disease
- if you are pregnant or breastfeeding your baby (see **Pregnancy and breastfeeding**)
- if you suffer from asthma and have previously experienced side effects to antihistamines
- Children under the age of two years should not be given **TEXA ALLERGY SYRUP**, as safety and efficacy have not been demonstrated.

Take special care with TEXA ALLERGY SYRUP:

- **TEXA ALLERGY SYRUP** is a non-sedating medicine, however some patients may become drowsy after taking the syrup (see **Driving and using machinery**)
- when taking **TEXA ALLERGY SYRUP** alcohol should not be consumed (see **Taking TEXA ALLERGY SYRUP with food and drink**)
- if you have any condition that results in urinary retention (difficulty in emptying your bladder)

PATIENT INFORMATION LEAFLET

TEXA ALLERGY SYRUP

- if you are an epileptic patient (have fits), or a patient at risk of convulsions
- if you are scheduled for allergy testing, you should stop taking **TEXA ALLERGY SYRUP** at least 3 days before testing, since it may affect your allergy test results
- if you are elderly, you are more likely to experience side effects of **TEXA ALLERGY SYRUP**.

Taking TEXA ALLERGY SYRUP with food and drink:

TEXA ALLERGY SYRUP can be taken with or without meals. Alcohol should be avoided.

Pregnancy and breastfeeding:

Do not use **TEXA ALLERGY SYRUP** if you are pregnant, planning to become pregnant or you are breastfeeding your baby (see **Do not take TEXA ALLERGY SYRUP**).

If you are pregnant or breastfeeding your baby, please consult your healthcare provider for advice before taking **TEXA ALLERGY SYRUP**.

Driving and using machinery:

TEXA ALLERGY SYRUP can cause drowsiness and you should exercise caution before driving, operating hazardous machinery or performing hazardous tasks.

Important information about some of the ingredients of TEXA ALLERGY SYRUP:

- **TEXA ALLERGY SYRUP** contains sorbitol and may have a laxative effect. If you have been told that you have an intolerance to some sugars, you should not take **TEXA ALLERGY SYRUP**.
- **TEXA ALLERGY SYRUP** contains the preservatives methyl parahydroxybenzoate and propyl hydroxybenzoate which may cause allergic reactions.

Taking other medicines with TEXA ALLERGY SYRUP:

Always tell your healthcare provider if you are taking any other medicine. (This includes

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

complementary or traditional medicines.)

- Taking any other medicines that may make you drowsy (sedatives, tranquilisers or strong painkillers) with **TEXA ALLERGY SYRUP** is not advised.

4. HOW TO TAKE TEXA ALLERGY SYRUP:

Do not share medicines prescribed for you with any other person. Always take **TEXA ALLERGY SYRUP** exactly as instructed in this leaflet. You should check with your doctor or pharmacist if you are not sure.

Adults or children 12 years of age or older:

10 mg (10 ml) once daily.

Children 6 to 12 years old:

10 mg (10 ml) daily, either as a single dose or as divided doses of 5 ml in the morning and 5 ml in the evening.

Children age 2 to 6 years:

5 mg (5 ml) daily, either as a single dose or as divided doses of 2,5 ml in the morning and 2,5 ml in the evening.

If you have a kidney or liver disorder, you should take 5 mg (5 ml) once daily.

If you take more TEXA ALLERGY SYRUP than you should:

In the event of an overdose, contact your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre.

Symptoms of an overdose may include, confusion, diarrhoea, dizziness, headache, general unwell feeling, dilated pupils, restlessness, drowsiness, stunned state in which it is difficult to get a response, feeling sleepy or tired, agitated, skin rash, passing less urine than is normal for you, uncontrolled trembling or shaking, and rapid heart rate.

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

If you forget to take TEXA ALLERGY SYRUP:

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the missed dose and take **TEXA ALLERGY SYRUP** at the next regularly scheduled time. Do not take a double dose to make up for the forgotten individual doses.

Effects when treatment with TEXA ALLERGY SYRUP is stopped:

When stopping treatment with **TEXA ALLERGY SYRUP**, you may experience severe itching or raised bumps on the skin. If this occurs, restart treatment with **TEXA ALLERGY SYRUP**, and the side effects should stop.

5. POSSIBLE SIDE EFFECTS:

TEXA ALLERGY SYRUP can have side effects.

Not all side effects reported for **TEXA ALLERGY SYRUP** are included in this leaflet. Should your general health worsen, or if you experience any untoward effects while taking **TEXA ALLERGY SYRUP**, please consult your healthcare provider for advice.

If any of the following happens, stop taking **TEXA ALLERGY SYRUP** and tell your doctor immediately or go to the casualty department at your nearest hospital:

- swelling of the hands, feet, ankles, difficulty in breathing, with or without swelling of the face, lips, tongue and/or throat which may cause difficulty in swallowing
- fainting
- severe itching of the skin (with raised lumps), skin rash.

These are all very serious side effects. If you have them, you may have had a serious allergic reaction to **TEXA ALLERGY SYRUP**. You may need urgent medical attention or hospitalisation.

Tell your doctor immediately or go to the casualty department at your nearest hospital if you

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

notice any of the following:

- seizures (fits)
- fast or irregular heartbeat
- thoughts of killing yourself (suicidal ideation)
- yellowing of the skin and eyes (jaundice), nausea, stomach pain, fatigue and fever; signs of a liver problem.

These are all serious side effects. You may need urgent medical attention.

Tell your doctor if you notice any of the following:

Frequent side effects:

- sore throat, runny and itchy nose and sneezing.

Less frequent side effects:

- anaemia with symptoms such as fatigue, pale skin, abnormal blood test results
- bleeding and bruising easily
- feeling sleepy or drowsy, depression, confusion, aggression, agitation, hallucinations (seeing and hearing things that aren't there), inability to sleep (insomnia)
- twitching, uncontrolled trembling or shaking
- unusual tiredness or weakness, generally feeling unwell
- blurred or double vision, eye strain, difficulty focussing, rolling of the eyes
- vertigo (feeling unbalanced / spinning head), ringing or buzzing in the ears (tinnitus)
- low blood pressure (light-headedness, dizziness)
- a wheezing sound when you inhale, thickening of mucous (phlegm)
- diarrhoea, abdominal pain, bloating, excess gas (wind), nausea, constipation, dry mouth
- abnormal liver test results
- skin rashes or other skin conditions such as itchy or red skin, or sensitivity to light
- hair loss, sweating

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

- muscle pain
- passing less urine than normal
- pain or discomfort when urinating
- wetting the bed, especially by children at night
- weight gain.

Side effects of unknown frequency:

- increased appetite
- headache, dizziness
- anxiety, nervousness, memory loss
- nightmares
- sensations of tingling, burning, pricking (pins and needles), or numbness of the skin
- bad taste in the mouth
- tremors (uncontrolled shaking)
- rash with blisters containing pus
- joint pain.

If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.

6. STORING AND DISPOSING OF TEXA ALLERGY SYRUP:

Store all medicines out of reach of children.

Store at or below 25°C in a dry place.

Do not use after the expiry date stated on the bottle.

Return all unused medicine to your pharmacist.

Do not dispose of unused medicine in drains or sewerage systems (e.g. toilets).

7. PRESENTATION OF TEXA ALLERGY SYRUP:

Type 3 amber glass bottle with a tamper evident, white coloured polyethylene closure,

PATIENT INFORMATION LEAFLET
TEXA ALLERGY SYRUP

contained in a printed outer carton. Pack sizes: 50 ml*, 100 ml, 150 ml* and 200 ml*.

*Not all pack sizes are marketed.

8. IDENTIFICATION OF TEXA ALLERGY SYRUP:

Clear or almost clear colourless solution with taste and odour of banana.

9. REGISTRATION NUMBER:

A41/5.7.1/0086

**10. NAME, BUSINESS ADDRESS AND TELEPHONE NUMBER OF THE REGISTRATION
HOLDER:**

Pharma Dynamics (Pty) Ltd

1ST Floor, Grapevine House, Steenberg Office Park

Silverwood Close

Westlake, Cape Town

7945, South Africa

Tel: + 27 21 707 7000

11. DATE OF PUBLICATION:

Date of registration: 08 February 2008

Date of latest approval: 07 January 2020

NAMIBIA: NS1 10/5.7.1/0332

ZAMBIA: P 051/004

MOZAMBIQUE: 2485